arXiv:1408.4292v1 [astro-ph.SR] 19 Aug 2014

Dwarf Nova EZ Lyncis Second Visit to Instability Strip

Elena P. Pavlenko,[1*,2] Taichi Kato,[2]
Aleksei A. Sosnovskij,[1] Maksim V. Andreev,[3,4] Tomohito Ohshima,[2]
Aleksander S. Sklyanov,[5] Ilfan F .Bikmaev,[5] Almaz I. Galeev,[5]

[1] *Crimean Astrophysical Observatory, 98409, Nauchny 19/17, Crimea, Ukraine*

[2] *Department of Astronomy, Kyoto University, Kyoto 606-8502*

[*] *eppavlenko@gmail.com*

[3] *Institute of Astronomy, Russian Academy of Sciences, 361605 Peak Terskol, Kabardino-Balkaria, Russia*

[4] *International Center for Astronomical, Medical and Ecological Research of NASU, Ukraine 27 Akademika Zabolotnoho Str. 03680 Kyiv, Ukraine*

[5] *Kazan Federal University, Kremlevskaya str.18, Kazan 420008, Russia*

(Received 201 0; accepted 201 0)

Abstract

The analysis of 14 periodograms of EZ Lyn for the data spaced over 565 d in 2012–2014 (2–3.5 yr after 2010 outburst) yielded the existence of the stable signals around 100 c/d and three signals around 310 c/d, 338 c/d and 368 c/d (the corresponding periods are 864 s, 279 s, 256 s and 235 s). We interpret them as independent non-radial pulsations of the white dwarf in EZ Lyn, but a possibility that a linear combination of frequency at 100 c/d and harmonic of orbital period could produce the frequency at 368 c/d also cannot be excluded. The signal at 100 c/d was detected during the first stay in the instability strip as a transient one. The period at 338 c/d, is a known non-radial pulsation EZ Lyn entered the instability strip after the 2010 outburst. We detected the signals around 310 c/d and 368 c/d for the first time. We applied the two-dimensional least absolute shrinkage and selection operator (Lasso) analysis for the first time to explore the behavior of these signals on the scale of hours for nightly runs of observations having duration of 6–12 hr. The Lasso analysis revealed the simultaneous existence of all three frequencies (310 c/d, 338 c/d and 368 c/d) for majority of nights of observations, but with variable amplitudes and variable drifts of frequencies by 2–6 percents on a time scale of ∼5–7 hr. The largest drift we detected corresponded to 17.5 s in period in ∼5 hours.

Key words: Accretion, accretion disks, — stars: novae, cataclysmic variables — stars: oscillations — stars: dwarf novae — stars: individual (EZ Lyncis)

1. Introduction

Cataclysmic variable (CV) is a close binary system containing a late-type component transferring matter via the inner Lagrangian point onto the primary component that is white dwarf. Due to an angular momentum loss, the CV with hydrogen-dominant atmosphere evolves to an orbital period minimum of 76.2 min (Knigge 2006) with decreasing of mass transfer rate up to 10^{-11} solar mass/year [Howell et al. (1995); Kolb, Baraffe (1999)]. The last circumstance enables the white dwarf to dominate the total radiation of CV, so such short-period systems are attractive for studies of the white dwarf, in particular to search for the white dwarf pulsations. Indeed, since 1998 non-radial pulsations of 14 accreting white dwarfs were found among the cataclysmic variables. They are GW Lib (Warner, van Zyl 1998), PQ And (Vanlandingham et al. 2005); V455 And (Araujo-Betancor et al. 2005); SDSS J133941.11+484727.5 = V355 UMa (Gänsicke et al. 2006); SDSS J091945.11+085710.1 (Mukadam et al. 2007); SDSS J151413.72+454911.9 = PP Boo (Nilsson et al. 2006); SDSS J074531.91+453829.5 = EQ Lyn (Mukadam et al. 2007); RE J1255+266 (Patterson et al. 2005); SDSS J150722.33+523039.8 = OV Boo (Patterson et al. 2008; Szkody et al. 2010); EZ Lyn (Pavlenko 2009); SDSS J161033.64−010223.3 = V386 Ser (Woudt, Warner 2004); SDSS J145758.21+514807.9 and BW Scl (Uthas et al. 2012). With two exceptions (OV Boo and RE J1255+266), their orbital periods are within the region of 80–90 min.

Knowledge of WD pulsations is important for measuring of its stellar mass, core composition, age, rotation rate, magnetic field strength, and distance (Winget, Kepler 2008; Fontaine, Brassard 2008).

EZ Lyn was discovered by Szkody et al. (2006) as a short period (0.059 d) quiescent dwarf nova with an underlying white dwarf. Pavlenko et al. (2007) first found it at the superoutburst. The large amplitude and series of 11 rebrightenings implied that this star belongs to the WZ Sge-type subclass of dwarf novae. After the outburst, Zharikov et al. (2008) found series of mini-outbursts with an amplitude about 0.5 mag.

In 2010 EZ Lyn had experienced the next superoutburst, which was discovered by H. Maehara (Kato et al. 2012). Contrary to the 2006 superoutburst, the 2010 superoutburst had six rebrightenings that is almost two times less than in the 2006 superoutburst. One month after the end of the main outburst, EZ Lyn was ∼1 magnitude fainter in the 2010 outburst than in the 2006 one.

The superhump period of EZ Lyn is 0.060 d [Shears et al. (2007), Pavlenko et al. (2007), Kato et al. (2009)], and its orbital period is 0.059005 d (Kato et al. 2009). Kato et al. (2009) first clarified the grazing eclipse of EZ Lyn during the 2006 superoutburst.

Eight months after the 2006 outburst, Pavlenko (2009) discovered the 756 s (12.6 min) brightness modulation that was interpreted as the nonradial pulsation of the white dwarf. This pulsation has been regularly detected and displayed a drift from 732 s to 768 s on a scale of

∼900 d (Pavlenko et al. 2012), and this wandering did not depend on time. Besides this stable dominating period, other periodic signals have appeared time by time between 846 s and 1302 s (Pavlenko et al. 2012).

Seven months after the 2010 outburst, EZ Lyn entered the instability strip the second time, but with a new 256–257 s pulsation (Pavlenko et al. 2012, Szkody et al. 2013). The drift of pulsation period as well as the change of pulsation amplitudes appears to be a common feature of accreting pulsators (Uthas et al. 2012; Mukadam et al. 2010). However, it is not clear what could be the shortest time scale of pulsation drift and change of its amplitude.

Here, during the second stay of EZ Lyn in the instability strip we present the detailed study of its periodic signals that are shorter then orbital period. We used least absolute shrinkage and selection operator (Lasso) analysis (Tibshirani 1996; Kato, Uemura 2012) for the first time to study the behavior of such signals on a time scale of hours.

2. Observations and data reduction

Observations of EZ Lyn have been carried out in the Crimean Astrophysical Observatory, the Terskol Observatory and using the Russian-Turkish 1.5-m telescope (RTT150) at TUBITAK National Observatory (Turkey) in 2012–2014 during 14 nights. The standard aperture photometry (de-biasing, dark subtraction and flat-fielding) was used for measuring of the EZ Lyn and comparison stars. We used the comparison stars the same as Zharikov et al. (2008). The observations have been carried out without usage of filters. Typical accuracy for the Crimean and Terskol observations was 0.005–0.017 mag and those of TUBITAK data was about 0.025 mag. The log of observation is given in table 1. Before the analysis we subtracted the smooth long-time trend for each night by using locally-weighted polynomial regression (LOWESS, Cleveland 1979). The times of observations are expressed in Barycentric Julian Dates (BJD). We used a phase dispersion minimization (PDM; Stellingwerf 1978), whose 1σ errors was estimated by the methods of Fernie (1989) and Kato et al. (2010); fast Fourier transform (FFT) [ISDA package, Pel't (1980)]. The false-alarm probability (FAP) was estimated according to Scargle (1982).

Lasso was introduced for period analysis of variable stars first time by Kato, Uemura (2012), which yielded very sharp signals. A two-dimensional Lasso power spectrum was found to be very effective in detecting varying multiple signals in Kepler observations of CVs [Kato, Maehara (2013); Osaki, Kato (2013); Kato, Osaki (2013)]. This method has been applied to the unevenly sampled ground-based data (Kato et al. 2014; Ohshima et al. 2014), and the resultant two-dimensional Lasso power spectra have been proven to have high frequency resolutions and are less affected by aliasing in unevenly sampled data than the conventional Fourier analysis.

Table 1. Log of the observations

Time BJD 2456000+	Telescope/CCD*	Exposure (s)	$N^{\dagger}$
236.360–236.619	A	60	242
237.341–237.488	A	45	181
244.518–244.626	A	60	120
245.336–245.623	A	45	415
247.340–247.627	A	60	304
248.388–248.627	A	90	202
249.333–249.488	B	30	246
250.427–250.661	B	20	716
306.238–306.579	B	30	579
307.156–307.672	B	20	1785
385.208–385.259	B	15	259
697.184–697.676	B	30	738
711.407–711.547	C	60	200
801.298–801.370	B	30	159

*A: 2 m Terskol/FLI PL430;

B: 2.6 m ZTSh/Apogee E47;

C: 1.5 m RTT-150/TFOSC

$^{\dagger}N$: Number of images

3. Periods

The complexity in the analysis of potential pulsations of the white dwarf in EZ Lyn is that they are contaminated by the orbital modulation. Both signals may be of variable amplitudes [Szkody et al. (2006), Pavlenko (2009)], but typically the orbital signal is the dominating one. The PDM spectrum and the mean orbital light curve (affected by the less-amplitude pulsations) for the data of 2013 January 14 are presented in figure 1. The strong first harmonic of the orbital signal is caused by the two-humped profile of the light curve. Its average amplitude is 0.08 mag, a grazing eclipse at the phase 0.7 is clearly seen. For this pattern the two humps are of slightly unequal heights, the smaller amplitude hump preceding the eclipse, has a small depression. This "splitting" of one or both humps could be sometimes more prominent, the periodograms of EZ Lyn often display the peak at 1/4 of orbital period (Pavlenko 2009, Kato et al. 2009).

For the data of every night we subtracted the orbital modulation with its first harmonic (that is consistent with two-humped curve) and calculated the power spectra. The result is given in figure 2.



Fig. 1. Above: PDM for 2013 January 14 (BJD 2456307). Below: phase-averaged data folded on the orbital 0.0590048 d period. The zero-epoch is BJD 2456307.16123. For clarity data are reproduced twice.

All the spectra contain indications to the periodic signals both at the low and high frequency regions. The signal around frequency 100 c/d (864 s) is surely seen at every night (the exception is BJD 2456237, when the significance of this signal was much lower comparing with those for other nights). The signal around 1/4 of orbital period is visible during all nights. The signal at 756 s being the dominating pulsation after the 2006 outburst, which was apparently a transient one and it is detected surely only on BJD 2456306, 2456307 and 2456697.

Meanwhile the behavior of the high-frequency signals is more complex and needs a detailed analysis.

The first power spectrum in this figure corresponds to the observations obtained ∼2 yr since the 2010 outburst of EZ Lyn (BJD 2456236). The 256-s pulsation that was detected by Pavlenko et al. (2012) seven months and Szkody et al. (2013) twelve months after the 2010 outburst, is the prominent and dominating one in a high-frequency part of the periodogram. But the next night the spectrum changed dramatically. The 256-s pulsation disappeared (or at least its amplitude reduced to the noise level). Instead one could see the marginal signals at 279 s and 235 s.

One week later (starting from BJD 2456244) the strong 235-s signal was detected. It



Fig. 2. The FFT for the nightly data of EZ Lyn in 2012–2014. The numbers in each frame denote the last three digits of the JD. The positions of dominating pulsation at 864 s and 235 s are shown by solid lines, and those at 756 s, 279 s and 256 s are shown by dotted lines. The horizontal line denotes the 0.1% FAP level.

remained to be the dominating one at least during the following week (BJD 2456244–2456250). At the same time starting from BJD 2456248 the signal at 279 s became prominent and its amplitude has increased up to BJD 2456248. Again this signal dropped practically to a noise level on BJD 2456306 and recovered on BJD 2456307. Since this date, the 279-s signal was registered as the dominating one in the high-frequency region. The signal at 256 s was detected close to the 0.1% FAP only on BJD 2456245 and BJD 2456250.

We suggest that all pulsations at the frequencies 100 c/d, 310 c/d, 338 c/d, 368 c/d (the corresponding periods are 864 s, 279 s, 256 s and 235 s) could be independent modes. However,



Fig. 3. Above: Light curve for BJD B2456244–2456250. Below: Lasso analysis ($\log\lambda = 2.5$). The width of the sliding window and the time step used are 0.15 d and 0.05 d, respectively.

there is an apparent linear combination of frequencies at 100 c/d (F_{100}), 368 c/d (F_{368}) and harmonic of the orbital period $F_{\rm orb}$: $F_{368} = 3F_{100} + 4F_{\rm orb}$. So this interpretation also cannot be excluded.

The similar problem was with identification of several pulsations in V386 Ser. While Woudt, Warner (2004) identified some pulsation as independent one, Mukadam et al. (2010) argued that it could be caused by a linear combination of orbital frequency and frequency of other known pulsation.

4. Two-dimensional Lasso analysis

We calculated the two-dimensional Lasso analysis for each of 14 nights separately in the region of frequencies around 100 c/d and in the region around high frequencies 275–400 c/d.

In figure 3 the Lasso spectra are shown for five closely spaced nights in 2012 November. It seems that these frequency displays a drift both from night to night and even within a night. More detailed behavior of this frequency is shown for the longest nightly data on BJD 2456307 (figure 4. This frequency displays a smooth quasi-periodic wandering within 98–100 c/d (or within $\sim$ 17 s) with typical time $\sim$4 hr.

To study the periods in the high frequencies region, we selected the patterns for the longest time series (6–12 hours) on BJD 2456236, 2456245, 2456247, 2456248, 2456250, 2456307, 2456697 and presented them in the figures 5, 6, 7, 8, 9, 10, 11. After consideration of the two-



Fig. 4. Above: Light curve for BJD 2456307. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.

dimensional power spectra it became evident that they give information on the behavior of detected periods over several hours. First, there is indication of the simultaneous co-existence of all three periods around 279 s, 256 s and 235 s (corresponding frequencies 310 c/d, 338 c/d and 368 c/d) for majority of nights. Second, the domination of these periods could change within several hours due to the non-synchronous increase or decrease of their amplitudes and some drift of period.

The data on BJD 2456236 show that the amplitude of the signal at the dominating frequency around 338 c/d was higher during the initial several hours. There is no clear indication of a signal at 368 c/d and there is some indication of a weak signal at 310 c/d of variable amplitude and phase. The signal at the dominating frequency around 368 c/d looks stable in its amplitude and period on BJD 2456245. There is a weaker but prominent signal at 338 c/d, which is stronger during the first hours of the observation. The frequency of this signal displays some decrease with time. The signal at frequency around 368 c/d for BJD 2456247, 2456248



Fig. 5. Above: Light curve for BJD 2456236. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.



Fig. 6. Above: Light curve for BJD 2456245. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.



Fig. 7. Above: Light curve for BJD 2456247. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.



Fig. 8. Above: Light curve for BJD 2456248. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.



Fig. 9. Above: Light curve for BJD 2456250. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.

and 2456250 remains to be the dominating one. The signal at frequency 338 c/d is rather marginal on BJD 2456247 and BJD 2456248, but is strong on BJD 2456250. The signal at frequency 310 c/d is strong for all these nights.

The longest 12-hr data set on BJD 2456307 revealed a signal at 310 c/d of variable amplitude and near-constant period that produces the dominating peak on the FFT periodogram. The most unusual thing is the behavior of a rather strong signal that varies between frequencies 340 c/d–352 c/d (this variation corresponds to 9 s in period) quasi-periodically on a typical time scale of ∼7 hr. This causes a "smeared" peak on the FFT periodogram. We believe this signal is related to those at frequency 338 c/d. Using all the data set, this signal varied by more than 9 s in period. There was no strong indication to the signal at 368 c/d.

The two-dimensional power spectrum on BJD 2456697 only shows a signal at 310 c/d with highly variable amplitude and increasing frequency from ∼305 c/d to ∼325 c/d that corresponds to the drift of 17.5 s in period in ∼5 hours.

5. Summary

The analysis of 14 periodograms of EZ Lyn for the data spaced over 565 d in 2012–2014 (2–3.5 yr after 2010 outburst) yielded the existence of several signals and their evolution. These are signals at the frequencies 100 c/d, 310 c/d, 338 c/d, 368 c/d (the corresponding periods are 864 s, 279 s, 256 s and 235 s). We believe that these signals arise from non-radial pulsations in



Fig. 10. Above: Light curve for BJD 2456307. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.



Fig. 11. Above: Light curve for BJD 2456697. Below: Lasso analysis ($\log\lambda = -2.8$). The width of the sliding window and the time step used are 0.15 d and 0.005 d, respectively.

the accreting white dwarf.

The transient appearance of pulsation at 846 s was detected during the first stay of EZ Lyn in the instability strip (Pavlenko 2009, Pavlenko et al. 2012), and the pulsation at 256 s was the pulsation when EZ Lyn entered the instability strip the second time. We detected the pulsations at 279 and 235 s for the first time. We suggest that all pulsations could be the independent modes, however a linear combination of frequencies at 100 c/d, 368 c/d and harmonic of the orbital period also cannot be excluded.

Our knowledge of a fast change in pulsation period and amplitude before this work had been rather limited. Recently Uthas et al. (2012) reported that pulsations in accreting white dwarfs BW Scl and SDSS J145758.21+514807.9 change in frequency by a few percent on a timescale of weeks or less; Chote, Sullivan (2013) found that in GW Lib a strong non-sinusoidal pulsation with a 19 minute period varied slightly in frequency over the six nights of observations.

Here we showed that the pulsations of EZ Lyn change in frequency by 2–6 percents on a time scale of hours and their amplitude may change on the same time scale. We found that Lasso is very powerful in detecting multiple frequencies and their variations in the pulsating white dwarfs.

We thank the anonymous referee whose comments greatly improved the paper. This work was supported by the Grant-in-Aid Initiative for High-Dimensional Data-Driven Science through Deepening of Sparse Modeling from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. The work is partially performed according to the Russian Government Program of Competitive Growth of Kazan Federal University; EP thanks to V. Malanushenko for valuable comments; SA, BI, GA thank to TUBITAK and KFU for partial support in using RTT150 (Russian-Turkish 1.5-m telescope in Antalya).

References

Araujo-Betancor, S., et al. 2005, A&A, 430, 629

Chote, P., & Sullivan, D. J. 2013, in ASP Conf. Ser. 469, 18th European White Dwarf Workshop, ed. J. Krzesinski, G. Stachowski, P. Moskalik, & K. Bajan (San Francisco: ASP), p. 337

Fernie, J. D. 1989, PASP, 101, 225

Fontaine, G., & Brassard, P. 2008, PASP, 120, 1043

Gänsicke, B. T., et al. 2006, MNRAS, 365, 969

Howell, S. B., Szkody, P., & Cannizzo, J. K. 1995, ApJ, 439, 337

Kato, T., et al. 2014, PASJ, 66, 30

Kato, T., & Maehara, H. 2013, PASJ, 65, 76

Kato, T., et al. 2012, PASJ, 64, 21

Kato, T., et al. 2010, PASJ, 62, 1525

Kato, T., & Osaki, Y. 2013, PASJ, 65, 97

Kato, T., et al. 2009, PASJ, 61, 601
Kato, T., & Uemura, M. 2012, PASJ, 64, 122
Knigge, C. 2006, MNRAS, 373, 484
Kolb, U., & Baraffe, I. 1999, MNRAS, 309, 1034
Mukadam, A. S., Gänsicke, B. T., Szkody, P., Aungwerojwit, A., Howell, S. B., Fraser, O. J., & Silvestri, N. M. 2007, ApJ, 667, 433
Mukadam, A. S., et al. 2010, ApJ, 714, 1702
Nilsson, R., Uthas, H., Ytre-Eide, M., Solheim, J.-E., & Warner, B. 2006, MNRAS, 370, L56
Ohshima, T., et al. 2014, PASJ, in press (arXiv/1402.5747)
Osaki, Y., & Kato, T. 2013, PASJ, 65, 95
Patterson, J., Thorstensen, J. R., & Kemp, J. 2005, PASP, 117, 427
Patterson, J., Thorstensen, J. R., & Knigge, C. 2008, PASP, 120, 510
Pavlenko, E. 2009, J. of Physics Conference Series, 172, 012071
Pavlenko, E., et al. 2012, Mem. Soc. Astron. Ital., 83, 520
Pavlenko, E., et al. 2007, in ASP Conf. Ser. 372, 15th European Workshop on White Dwarfs, ed. R. Napiwotzki, & M. R. Burleigh (San Francisco: ASP), p. 511
Pel't, Ya. 1980, Frequency analysis of astronomical time series. (Tallin: Valgus)
Scargle, J. D. 1982, ApJ, 263, 835
Shears, J., Klingenberg, G., & de Ponthiere, P. 2007, J. Br. Astron. Assoc., 117, 331
Stellingwerf, R. F. 1978, ApJ, 224, 953
Szkody, P., et al. 2006, AJ, 131, 973
Szkody, P., et al. 2010, ApJ, 710, 64
Szkody, P., Mukadam, A. S., Sion, E. M., Gansicke, B. T., Henden, A., & Townsley, D. 2013, AJ, 145, 121
Tibshirani, R. 1996, J. R. Statistical Soc. Ser. B, 58, 267
Uthas, H., et al. 2012, MNRAS, 420, 379
Vanlandingham, K. M., Schwarz, G. J., & Howell, S. B. 2005, PASP, 117, 928
Warner, B., & van Zyl, L. 1998, in 185, New Eyes to See Inside the Sun and Stars, ed. F.-L. Deubner, J. Christensen-Dalsgaard, & D. Kurtz (Berlin: Springer), p. 321
Winget, D. E., & Kepler, S. O. 2008, ARA&A, 46, 157
Woudt, P. A., & Warner, B. 2004, MNRAS, 348, 599
Zharikov, S. V., et al. 2008, A&A, 486, 505